Original Research Article

Histopathological study of lesions of nasal cavity and paranasal sinuses

Arpana Kulkarni¹, Abhinav Shetty^{1,*}, Pooja Pathak¹¹Dept. of Pathology, Dr. D. Y. Patil Medical College, Hospital and Research Centre, Dr. D. Y. Patil Vidyapeeth, Pimpri, Pune, Maharashtra, India

ARTICLE INFO

Article history:

Received 01-06-2019

Accepted 29-08-2019

Available online 22-02-2020

Keywords:

Nasal cavity

Paranasal sinuses

Sinonasal polyp

Sinonasal papilloma

ABSTRACT

Introduction: The nasal cavity and paranasal sinuses are exposed to many environmental pollutants and pathogens. Exposure to such influences can lead to a variety of lesions primarily affecting the sinonasal tract. The aim of this study was to find out the incidence, age and sex distribution and to enlist the different types of lesions of the nasal cavity and paranasal sinuses.

Materials and Methods: A study of 122 cases was conducted over a 2 year period. Both retrospective and prospective cases were included in the study. All relevant clinical details, tissue sections with H&E, special stains and IHC stains were done whenever necessary.

Result: Out of 122 cases, 98 non-neoplastic cases and 24 neoplastic cases were diagnosed. Maximum non-neoplastic cases were detected in the third and fourth decade. Benign lesions were commonly noted in fourth and seventh decade. Malignant lesions of sinonasal tract commonly afflicted patients of 61 to 70 years of age. Non-neoplastic and benign lesions showed male predominance. Female preponderance was noted in malignant lesions of sinonasal tract. Among non-neoplastic lesions, sinonasal polyp (84 cases) was most prevalent. Maximum proportion of benign lesions were diagnosed as hemangioma (6 cases) and sinonasal papilloma (5 cases). Among malignant neoplastic lesions, 4 cases were of squamous cell carcinoma and 3 cases were of malignant melanoma.

Conclusion: A variety of lesions with overlapping clinical features can affect the sinonasal tract. Histopathology remains the gold standard for establishing the diagnosis in such cases.

© 2020 Published by Innovative Publication. This is an open access article under the CC BY-NC-ND license (<https://creativecommons.org/licenses/by/4.0/>)

1. Introduction

'Sinonasal tract' is a collective term used for nasal cavity and paranasal sinuses.¹ The main functions of the sinonasal tract are filtering and humidifying inhaled air. The nasal cavity also has specific olfactory receptors for airborne odorant molecules. The paranasal sinuses additionally have the functions of acting as resonating chambers during speech and reducing the relative weight of the skull.²

These functions lead to exposure of various allergens, pathogens, chemical and physical irritants and other environmental influences. As a result of these multifaceted exposures, various inflammatory conditions, infections and neoplasms can occur in the sinonasal tract.³

Prevalence of sinonasal masses in the general population is 1 to 4%.⁴ A careful clinical workup including symptomatology, radiological investigations and endoscopy helps to determine a differential diagnosis but histopathology provides the final diagnosis.⁵ Thus histopathological examination is essential for timely diagnosis and intervention.⁶

2. Material and Methods

This study was a descriptive cross-sectional study and was carried out in the Pathology department of a tertiary care hospital. Institute Ethics Committee Clearance (IECC) was obtained before start of the study.

The study took 2 years for completion and 122 cases from September 2011 to August 2018 were studied. Cases from September 2011 to August 2016 were part of the

* Corresponding author.

E-mail address: abhinav320@gmail.com (A. Shetty).

retrospective study. The archives of the Department of Pathology provided the paraffin blocks, slides and reports required for the retrospective study. For the prospective study, the formalin fixed tissue specimens were received with a detailed clinical history and radiological findings. Gross findings of the specimens was noted and adequate sections were taken and stained with hematoxylin and eosin. Special stains and immunohistochemical stains were used to confirm diagnosis wherever necessary. Neoplastic lesions diagnosed were classified accordingly by referring the 2017 WHO classification of tumours of nasal cavity and paranasal sinuses.⁷

3. Result

Lesions of the sinonasal tract commonly affected the age group 31-40 years (28 cases, 22.95%) followed by 21-30 years (27 cases, 22.13%). Maximum patients of non-neoplastic cases belonged to 21-30 and 31-40 years age groups (24 cases each, 24.49%). Benign lesions were mostly noted in fourth and seventh decade of life (3 cases each, 23.08%) whereas a high proportion of malignant lesions were reported in 61-70 years age group (4 cases, 36.36%). (Table 1)

Sinonasal tract lesions were commonly seen in males. 71 out of 122 cases (58.20%) were males and 51 cases (41.80%) were females. Out of the 98 non-neoplastic cases diagnosed, majority were males (61 cases, 62.25%) whereas female preponderance was noted among neoplastic lesions (14 of 24 cases, 58.33%).

Of the total 122 cases included in this study, the non-neoplastic cases (98 cases) outnumbered the neoplastic cases (24 cases). Cases diagnosed as benign lesions (13 cases, 10.66%) were marginally higher than those diagnosed as malignant (11 cases, 9.01%).

In terms of frequency, the non-neoplastic lesions of sinonasal tract diagnosed were, in descending order, sinonasal polyp, fungal rhinosinusitis, epidermal inclusion cyst, rhinosporidiosis, chronic non-specific inflammation, lepromatous leprosy and arteriovenous malformation (Table 2). Benign lesions comprised of hemangioma, inverted sinonasal papilloma, angiofibroma and trichoeplithelioma (Table 3). Squamous cell carcinoma, malignant melanoma, nasopharyngeal carcinoma, adenoid cystic carcinoma and T-cell lymphoma were the malignant lesions noted (Table 4).

4. Discussion

In the present study, age of presentation showed a wide range from 2 to 90 years. Maximum cases were noted in the fourth decade (28 cases, 22.95%) followed by the third decade (27 cases, 22.13%). Raj et al found maximum cases in third decade (32.79%) followed by fourth decade (21.31%).⁸ In the study conducted by Mane et al, the most

Fig. 1: Photomicrograph of sinonasal polyp. Ciliated pseudostratified columnar respiratory epithelium with stromal mixed inflammatory infiltrate. (H&E, 400X)

Fig. 2: Photomicrograph of aspergillus (mycetoma). Aggregates of hyphae noted. Hyphae are thin and branch at acute angles. (H&E, 400X) Inset: GMS stain highlights the thin, branching hyphae

common age group was 21-30 years (35 cases, 27.78%) followed by 31-40 years (31 cases, 24.60%).⁹

Male to female ratio of 1.39:1 was calculated in this study. Similar findings of male preponderance were observed in studies conducted by Tondon et al,¹⁰ Khan et al¹¹ and Kulkarni et al.⁶

We observed higher frequency of non-neoplastic lesions (80.33%) in comparison with neoplastic lesions (19.67%). Our findings are in concordance with Khan et al,¹¹ Kulkarni et al⁶ and Mane et al.⁹ However Dasgupta et al¹² recorded

Table 1: Age distribution of lesions of nasal cavity and paranasal sinuses

Age Group	Non-neoplastic	Neoplastic Benign	Malignant	No. of cases	% of total cases
0 to 10	4	0	0	4	3.28%
11 to 20	9	2	1	12	9.84%
21 to 30	24	2	1	27	22.13%
31 to 40	24	3	1	28	22.95%
41 to 50	13	2	3	18	14.75%
51 to 60	15	1	1	17	13.93%
61 to 70	6	3	4	13	10.66%
>70	3	0	0	3	2.46%
Total	98	13	11	122	100%

Table 2: List of non-neoplastic lesions of sinonasal tract diagnosed in the present study

Histopathological diagnosis	No of cases	Percentage (%)	% of total cases
Non-neoplastic lesions	98	100%	80.33%
1. Sino-nasal Polyps	84	85.72%	68.85%
2. Fungal rhinosinusitis	5	5.10%	4.10%
3. Epidermal inclusion cyst	4	4.08%	3.28%
4. Rhinosporidiosis	2	2.04%	1.64%
5. Chronic non-specific inflammation	1	1.02%	0.82%
6. Lepromatous leprosy	1	1.02%	0.82%
7. Arteriovenous malformation	1	1.02%	0.82%

Table 3: List of benign neoplastic lesions of sinonasal tract detected in the present study

Histopathological diagnosis	No of cases	Percentage (%)	% of total cases
Benign neoplastic lesions	13	100%	10.66%
1. Hemangioma	6	46.16%	4.92%
2. Inverted sinonasal papilloma	5	38.46%	4.10%
3. Angiofibroma	1	7.69%	0.82%
4. Trichoepithelioma	1	7.69%	0.82%

Table 4: List of malignant neoplastic lesions of sinonasal tract detected in the present study

Histopathological diagnosis	No of cases	Percentage (%)	% of total cases
Malignant neoplastic lesions	11	100%	9.01%
1. Squamous cell carcinoma	4	36.37%	3.27%
2. Malignant melanoma	3	27.27%	2.46%
3. Nasopharyngeal carcinoma	2	18.18%	1.64%
4. Adenoid cystic carcinoma	1	9.09%	0.82%
5. T-cell lymphoma	1	9.09%	0.82%

an almost equal proportion of neoplastic and non-neoplastic lesions. (Table 5)

4.1. Non-neoplastic lesions of sinonasal tract

In the present study, majority of the lesions were of non-neoplastic etiology. The most common non-neoplastic lesion was sinonasal polyp (85.72%) followed by fungal rhinosinusitis (5.10%), epidermal inclusion cyst (4.08%), rhinosporidiosis (2.04%), lepromatous leprosy (1.02%), chronic nonspecific inflammation (1.02%) and arteriovenous malformation (1.02%). Studies conducted by Dasgupta et al,¹² Pushpalatha et al¹³ and Mane et al⁹

also noted that sinonasal polyp was the most common non-neoplastic lesion. Other non-neoplastic lesions like fungal rhinosinusitis, rhinosporidiosis and epidermal inclusion cyst showed varying frequencies.

4.2. Sinonasal polyps

Peak incidence of sinonasal polyps was noted in the third decade. Similar findings were observed by Khan et al¹¹ and Kulkarni et al⁶ who reported a peak incidence in second and third decade of life. Male predominance was observed and this was in compliance with the studies by Dasgupta et al¹² and Khan et al.¹¹ Microscopically, the epithelial lining

Table 5: Frequency of non-neoplastic and neoplastic lesions compared to previous studies

	Dasgupta et al ¹² (1997)	Khan et al ¹¹ (2006)	Kulkarni et al ⁶ (2012)	Mane et al ⁹ (2017)	Present study (2018)
Non-neoplastic lesions	50.70%	60%	86%	83.33%	80.33%
Neoplastic lesions	49.30%	40%	14%	16.67%	19.67%

Fig. 3: Photomicrograph of rhinosporidiosis. Subepithelial and submucosal sporangia containing sporangiospores noted. (H&E, 100X) **Fig. 5:** Photomicrograph of inverted sinonasal papilloma. Surface epithelium showing multiple inversion with loss of seromucinous glands in the submucosa. (H&E, 100X) **Fig. 4:** Photomicrograph of capillary hemangioma. Well circumscribed, lobular arrangement with capillary proliferation. Overlying collarette of epithelium is noted. (H&E, 100X) **Fig. 6:** Photomicrograph of malignant melanoma. Melanocytic atypia with prominent nucleoli and finely dispersed chromatin. (H&E, 400X) Inset: HMB-45 immunoreactivity noted. S-100 reactivity was also noted.

Fig. 7: Photomicrograph of Adenoid cystic carcinoma. Typical cribriform pattern. Nests of cells arranged concentrically around gland-like spaces. (H&E, 100X)

is ciliated pseudostratified columnar, with stromal infiltrate comprising of eosinophils, lymphocytes and neutrophils. (Figure 1)

4.3. Fungal rhinosinusitis

According to this study, the mean age of presentation of patients diagnosed with fungal rhinosinusitis was 62.4 years with female predominance (M:F ratio 1:1.5). Mane et al⁹ reported the mean age of presentation of fungal rhinosinusitis was 50.7 years with male preponderance (2:1). This suggests that fungal rhinosinusitis mainly affects the elderly. On microscopy, the fungal hyphae were thin and branching at acute angles. (Figure 2)

4.4. Neoplastic lesions of sinonasal tract

In the present study, benign neoplastic lesions (54.17%) marginally outnumbered the malignant neoplastic lesions (45.83%). Similar findings were noted in the study by Khan et al.¹¹ But Tondon et al¹⁰ and Mane et al⁹ reported a higher proportion of benign neoplastic lesions (73.53% and 71.43% respectively) compared to malignant lesions (26.47% and 28.57% respectively).

4.5. Benign neoplastic lesions of sinonasal tract

Maximum benign neoplastic lesions of sinonasal tract in our study were seen in the 31 to 40 years and 61 to 70 years age group. Mane et al⁹ reported similar findings where benign tumours were reported commonly in fourth decade. Male preponderance was reported in benign neoplastic lesions and this was corroborated by Kulkarni et al,⁶ Raj et al⁸ and

Mane et al.⁹

Hemangioma (46.16%) was the commonest benign lesion followed by inverted sinonasal papilloma (38.46%), angiofibroma (7.69%) and trichoepithelioma (7.69%). Kulkarni et al⁶ observed that the most common benign neoplastic lesion of sinonasal tract was hemangioma (38.46%) followed by angiofibroma (30.76%). Mane et al⁹ noted sinonasal papilloma (40%) and Khan et al¹¹ reported angiofibroma (42.85%) as the most common benign neoplastic lesion of sinonasal tract.

4.6. Hemangioma

It was noted that hemangioma could occur in any age group but was most commonly noted in the fourth decade. This finding was in accordance with Dasgupta et al¹² and Kulkarni et al⁶ who reported the same and the mean age of presentation in their studies was 32.4 years. Females were more afflicted than males (1:2). Similar findings were noted by Kulkarni et al⁶ (1:1.5) but Dasgupta et al¹² observed male preponderance (1.95:1). Microscopic features such as lobular arrangement, capillary proliferation and capillary lining by plump endothelial cells were noted. (Figure 4)

4.7. Inverted sinonasal papilloma

Mean age of presentation of patients diagnosed with inverted papilloma was 54.6 years which was in concordance with studies by Tondon et al¹⁰ and Mane et al.⁹ Inverted papilloma predominantly affected males in this study (M :F ratio is 4:1). Similar observations were noted by Tondon et al¹⁰ and Mane et al⁹ (M:F ratio 3:1 and 5:1 respectively). Microscopic examination revealed inverted growth pattern of surface epithelium with intact and distinct basement membrane and loss of submucosal glands. (Figure 5)

4.8. Malignant neoplasms of sinonasal tract

Maximum cases of malignant neoplasms of sinonasal tract were observed in the seventh decade of life in the present study. Khan et al¹¹ and Mane et al⁹ reported maximum malignant neoplasms in sixth and seventh decade. Malignant neoplasms were common in females in the present series (M :F ratio was 1:4.5). Similar findings were noted by Raj et al⁸ (1:1.67) and Mane et al⁹ (1:2).

The most common malignant neoplastic lesion of sinonasal tract was squamous cell carcinoma (36.37%) followed by malignant melanoma (27.27%), nasopharyngeal carcinoma (18.18%), adenoid cystic carcinoma (9.09%) and T cell lymphoma (9.09%). Studies by Khan et al¹¹ and Raj et al⁸ also reported squamous cell carcinoma as the most common malignant lesion but Mane et al⁹ reported sinonasal undifferentiated carcinoma as the most common malignant sinonasal tract lesion. Other malignant neoplastic lesions diagnosed in this study showed varying frequencies

or were not noted at all.

4.9. Squamous cell carcinoma

The age range of patients with squamous cell carcinoma observed was 18 to 65 years and the mean age of presentation calculated was 39.5 years. Dasgupta et al¹² noted the mean age of presentation as 52.3 years for patients suffering from sinonasal squamous cell carcinoma whereas Khan et al¹¹ reported peak incidence of squamous cell carcinoma in the sixth and seventh decade of life. We found that only females were affected by squamous cell carcinoma. However, Dasgupta et al¹² and Khan et al¹¹ noted a male predominance (2:1). This variation can be attributed to the small sample size (4 cases) in our study. On microscopy, infiltrating nests of squamous epithelial cells were noted. The epithelial cells were highly pleomorphic with hyperchromatic nuclei, nuclear atypia and abnormal mitotic figures. Keratin pearls were observed.

4.10. Malignant melanoma

Peak incidence of malignant melanoma was in the seventh decade. Khan et al¹¹ noted maximum cases in the fourth decade. In the current series, all malignant melanoma cases were females. Khan et al¹¹ noted equal sex distribution. Microscopic examination revealed atypia of melanocytes with prominent nucleoli and mitotic activity. On immunohistochemistry, neoplastic cells showed reactivity for HMB-45 and S-100. (Figure 6)

5. Conclusion

To conclude, various types of lesions affect the sinonasal tract. These lesions can affect any age group. Irrespective of their etiology, whether non-neoplastic or neoplastic, the lesions can present with similar clinical features. Radiological studies and other investigations can help provide a differential diagnosis, but the final diagnosis can only be given on histopathological examination. Definite diagnosis is essential for further management and to determine the prognosis of the patient. Therefore histopathology plays an integral role and remains the gold standard for establishing the diagnosis of sinonasal tract lesions.

6. Source of funding

None.

7. Conflict of interest

None.

References

1. Bhattacharyya R, Chakrabarti I, Giri I, Banerjee A, Bhattacharya J, Goswami B. A clinicopathological study of masses arising from sinonasal tract and nasopharynx in north Bengal population with special reference to neoplasms. *Egypt J Otolaryngol*. 2015;31(2):98.
2. Shah SN, Goswami Y. Study of Lesions of Nasal Cavity, Nasopharynx and Paranasal Sinuses by Histopathological Examination. *Gujarat Med J*. 2012;67(2):70–72.
3. Jaison J, Tekwani D. Histopathological lesions of nasal cavity, paranasal sinuses and nasopharynx. *Ann Appl Bio-Sci*. 2015;2(2):40–46.
4. Kirtsreesakul V. Update on Nasal Polyps: Etiopathogenesis. *J Med Assoc Thai*. 2005;88(12):1966–1972.
5. Zafar U, Khan N, Afroz N, Hasan S. Clinicopathological study of non-neoplastic lesions of nasal cavity and paranasal sinuses. *Indian J Pathol Microbiol*. 2008;51(1):26.
6. Kulkarni A, Mudholkar V, Acharya A, Ramteke R. Histopathological Study of Lesions of Nose and Paranasal Sinuses. *Indian J Otolaryngol Head Neck Surg*. 2012;64(3):275–279.
7. El-Naggar A, Chan J, Grandis J, Slootweg P, Takata T. WHO classification of head and neck tumours. 4th ed. and others, editor ; 2017.
8. Raj JA, Sharmila PS, Shrivastava M, Mahantachar V, Rajaram. Morphological spectrum of lesions in the sinonasal region. *J Evol Med Dent Sci*. 2013;2(37):7175–7186.
9. Mane PS, Agale SV. Clinicopathological Study of Sinonasal Masses. *Annals Pathol Laboratory Med*. 2017;4(3):261–267.
10. Tondon PL, Gulati J, Mehta N. Histological study of polypoidal lesions in the nasal cavity. *Indian J Otolaryngol Head Neck Surg*. 1971;13:3–11.
11. Khan N, Zafar U, Afroz N, Ahmad SS, Hasan SA. Masses of nasal cavity, paranasal sinuses and nasopharynx: A clinicopathological study. *Indian J Otolaryngol Head Neck Surg*. 2006;58(3):259–263.
12. Dasgupta A, Ghosh RN, Mukherjee C, & Hns I ; 1997.
13. Pushpalatha K, Ch, & R S, Soujanya. A study on Histomorphological Spectrum of Nasal Polyp. *Ann Pathol Lab Med*. 2017;4:153–156.

Author biography

Arpana Kulkarni Associate Professor

Abhinav Shetty Resident

Pooja Pathak Resident

Cite this article: Kulkarni A, Shetty A, Pathak P. Histopathological study of lesions of nasal cavity and paranasal sinuses. *Indian J Pathol Oncol* 2020;7(1):88-93.