Original Research Article

Mean platelet volume: Has it the capacity to predict the concentration of Platelet counts in Dengue: A study

Anagha A Joshi¹, Sathyavathi S^{2,*}, Gayathri B R¹, Rukmini Tanjore³, Krishnamurthy B N²

¹Dept. of Pathology, Kempegowda Institute of Medical Sciences, Bengaluru, Karnataka, India

²Dept. of Pathology, Sri Jayadeva Institute of Cardiovascular Sciences and Research, Bengaluru, Karnataka, India

³Prathima Institute of Medical Sciences, Karim Nagar, Telangana, India

ARTICLE INFO

Article history:

Received 16-03-2020

Accepted 24-04-2020

Available online 19-08-2020

Keywords:

MPV

Dengue

Platelet

ABSTRACT

Background: Dengue, endemic in India presents in mild to severe forms with or without bleeding. There is no specific treatment available till date hence severe dengue has to be recognized in early stages. Platelet counts are unreliable as they do not correlate with bleeding. Platelet indices have been studied for their utility as predictors of severity of the disease and aid in guide to transfusion therapy. Mean platelet volume is inexpensive and available with routine complete blood counts in hematology analyser. Its role as marker in diagnosis, prognosis and treatment of dengue has been investigated with platelet indices reports.

Aims: To study MPV patterns, association with platelet counts and relevant laboratory parameters in dengue.

Methods: The study was conducted over one month in the year 2016 in department of Hematology, Kempegowda Institute of Medical Sciences Hospital, Bengaluru. 100 serologically proven dengue cases were analysed with data collected from the records in the departments of Hematology and Microbiology against patient's unique hospital identification number.

Results: There was a predominance of young male patients. The range for MPV was 8.7 – 13.2 (average 10.95) with 35% normal and 65% high MPV cases. There was 48% cases with moderately severe / severe thrombocytopenia. 32% with increased hematocrit, 35% with leucopenia, 54% with significant atypical lymphocytosis and 42% with antibody pattern. There was an increased number of high MPV seen in severe thrombocytopenia, increased hematocrit significant atypical lymphocytosis and antibody pattern whereas normal pattern was noted in leucopenia.

Conclusion: High MPV is a useful predictor of severe dengue platelet recovery and may guide in transfusion therapy.

© 2020 Published by Innovative Publication. This is an open access article under the CC BY-NC license (<https://creativecommons.org/licenses/by-nc/4.0/>)

1. Introduction

Dengue of serious public health concern in India caused by DENV, manifests as asymptomatic mild or severe dengue with haemorrhage and shock.¹ Thrombocytopenia is one of the criteria to indicate the clinical severity and also contributes to bleeding. Platelet counts may or may not directly correlate with haemorrhagic manifestations.^{1–5} Platelet counts decrease from the 3rd to 7th day

and usually normalizes by the 8th to 10th day of fever. Thrombocytopenia is caused by the bone marrow suppression by the virus, increased peripheral destruction or increased consumption / abnormal pooling of platelets.^{6,7} Management dilemmas occur with severe thrombocytopenia due to risk of haemorrhage / bleeding leading to unwarranted, unsafe, excess and expensive transfusions.^{8–10}

A 'wait and watch' restrictive transfusion policy is beneficial. Laboratory tests like Platelet indices which aid in patient management have been investigated. Mean Platelet Volume (MPV) the volume of platelets measured in

* Corresponding author.

E-mail address: sathyavathikiran@gmail.com (Sathyavathi S).

femtolitres (fl) is a predictor of platelet recovery, etiology of thrombocytopenia (hypoproliferative or rapidly responding bone marrow), worsening dengue and impacts therapy.^{1,6,10}

2. Aims and Objectives

Our study investigated the association of MPV with

1. Platelet count and relevant laboratory parameters
2. Its patterns in Dengue.

3. Materials and Methods

This study included 100 serologically proven dengue cases at department of Haematology, Kempe Gowda Institute of medical sciences hospital, Bengaluru over a period of one month in the year 2016.

The Hematology data obtained by the Hematology Analyser Sysmex 1800i retrieved from the hematology records and the Dengue serology from the microbiology department records was tabulated. Platelet and Differential counts was checked on peripheral smear stained with Leshman stain as per the hospital protocol for confirmation of counts. All data was analysed.

3.1. Inclusion criteria

All serologically proven cases of Dengue with thrombocytopenia and complete hemogram data.

3.2. Exclusion criteria

Cases of Dengue with associated diseases, normal and increased platelet counts and without relevant lab data.

3.3. Ethical committee clearance

The patient data was analysed against unique hospital identification number and anonymity maintained. The study was approved by the Ethical committee of hospital.

4. Results

Our study showed an age range from 9 months to 65 years with a male to female ratio of 1.3:1 (Table 1).

The analysis of MPV patterns showed a range from 8.7 to 13.2 fl with an average of 10.95 fl. The normal MPV range for the study was 7.5 to 10.5 fl.

We had 35% to 65% of cases with normal and increased MPV respectively (Table 2).

Thrombocytopenia was graded as

- 1) Mild : 0.76 – 1.49 l/cumm in 28% of cases
- 2) Moderate : 0.51- 0.75 l/cumm in 24% cases
- 3) Moderately Severe : 0.26 – 0.50 l/cumm in 32% cases
- 4) Severe : < 0.25 l/cumm in 16% cases.

MPV association with Thrombocytopenia (Table 3)

Normal MPV was noted in 12/35 (34%) as against 36/65 (55%) with increased MPV in moderately severe thrombocytopenia. Normal MPV was seen in 6/16 (38%) as against 10/16 (62%) of high MPV cases in counts < 0.25 l/cumm. As the platelet counts increased, the proportion of both became approximately equal.

MPV with hematocrit (Table 4) showed 32% with hematocrit greater than 45, of which 69% had high and 31% had normal MPV. Analysing MPVs 22/65 (34%) of increased and 10/35 (28%) of normal MPV had high hematocrit.

MPV with Leucopenia (Table 5) showed 35% with Leucopenia, 55% with normal counts and 10% with Leucocytosis.

20/65 (31%) of increased as against 15/35 (43%) of normal MPV had leucopenia.

The Atypical Lymphocytosis is significant if >20% lymphocytes were found to be of atypical type. Of the 54 cases, there were 40/54 (74%) with significant increased atypical lymphocytosis as against 14/54 (26%) with normal MPV. Analysing MPV, 14/35 (40%) normal as against 40/65 (62%) with high MPV showed significant Atypical lymphocytosis.

We had 15/35 (43%) of normal cases as 15/65 (25%) of high MPV with NS1 antigen pattern. Antibody association was noted in 30/65 (46%) of high as against 12/35 (34%) of normal MPV. The distribution of NS1 antigen was equal between normal and high MPV but antibody pattern showed a predominance of high MPV.

5. Discussion

Platelet indices have been the focus of interest for their utility in Dengue.¹¹ The tests are simple, inexpensive and non invasive.^{10,12–16} MPV is an indicator of platelet size and is reported as a ratio of Plateletcrit to total platelet count in Hematology Analyser by different methods like Impedence, optical light scatter counting etc.^{10,11,16,17} Normal MPV range for our study is 7.5 – 10.5 fl^{13,16} but differ in other due to many variables.^{11,14,16–18} In healthy individuals, MPV correlates inversely with platelet counts but the proportions differ in pathologic states.¹⁷

MPV is indicative of platelet turnover and is increased with rapid marrow response and it decreases with hypoproliferative states in Dengue with thrombocytopenia^{6,10,12,19} because

1. The younger platelets are larger in size and the newly formed cells contain more granules.
2. Activated platelets change shape from biconcave to spheroidal with pseudopod formation.
3. In hypoproliferative states only aged, small platelets remain in circulation.^{10,13–16,18–21}

MPV correlates with platelet activation and recovery indicating the etiology of thrombocytopenia

Table 1: Age and sex pattern

Age				Sex				Total
>12 years (Adults)		< 12 years (Pediatric)		Male		Female		
n	%	N	%	n	%	n	%	
71	71	29	2	57	57	43	43	100

Table 2: MPV (fl) patterns

MPV Range	N	%
< 7.5	Normal	Normal
7.5 – 10.5	35	35
10.6 – 11.5	44	44
11.6 – 12.5	16	16
>12.6	05	05
Total	100	100

Table 3: MPV & Platelet counts

MPV	< 0.25		0.26 – 0.5		0.5 – 0.75		0.76 – 1.5		Total
	n	%	n	%	n	%	n	%	
< 7.5									
7.5 – 10.5	06	17	06	17	12	34	11	32	35
10.6 – 11.5	07	16	17	39	07	16	13	29	44
11.6 – 12.5	02	12	08	50	03	19	03	19	16
>12.6	01	20	01	20	02	40	01	20	05
Total	16		32		24				100

Table 4: MPV and Hematocrit

		Hematocrit				
MPV (fl)		< 45 %		>45 %		Total
	N	%	n	%		
7.5 – 10.5	25	71	10	29		35
>10.6	43	66	22	34		65
Total	68		32			

Table 5: MPV with total white cell count (cells / cumm)

MPV (fl)	Normal (4000 -11000)		Leucopenia (< 4000)		Leucocytosis (> 11000)		Total
	n	%	n	%	n	%	
7.5- 10.5	19	54	15	4	01	03	35
>10.6	36	55	20	31	09	14	65
Total	55		35		10		100

Table 6: Atypical Lymphocytosis

MPV (fl)	< 20 %		>20 %		Total
	n	%	n	%	
7.5 – 10.5	21	60	14	40	35
>10.6	25	38	40	62	65
Total	46		54		100

Table 7: MPV with Serology

MPV (fl)	NS 1 pattern		Mixed		Antibody Pattern		Total
	N	%	N	%	n	%	
7.5 – 10.5	15	43	08	23	12	34	35
>10.6	15	25	20	31	30	46	65
Total	30		28		42		100

(hyperdestructive due to peripheral destruction of platelets with rapid marrow turnover or hypoproliferative due to marrow suppression by virus) and impacts prognosis and treatment.^{6,11–13,15,16}

Our study shows a predominance of younger aged and male patients in accordance with the other studies.^{13,15,22}

MPV patterns showed a range of 8.7 – 13.2 but differed in others.^{6,13,15,21,23} The average MPV was in accordance with few studies like Basher et al & Pritam Sewakdar et al^{6,10,14} but varied in others like Gogaram et al^{19,24,25} MPV patterns were similar to the study by Ishito et al. thrombocytopenia patterns were in accordance with the same³ but varied in others.^{2,4}

Our study showed a high MPV in increased proportion of thrombocytopenia cases in accordance with few^{24,25} but differed in others²¹ we had a predominance of high MPV associated with severe thrombocytopenia and high hematocrit as per the WHO criteria for severe dengue. Increased hematocrit is attributed to the endothelial damage by cytokines which also cause platelet activation, increased size and MPV.^{26,27} We had 34% of high MPV in association with increased hematocrit and 60% cases with platelet count less than one lakh in accordance with study of K. Sharma et al with high MPV in 38.8% Dengue shock syndrome and 56.5% of Dengue haemorrhagic fever cases respectively. There was increased proportion of high MPV cases with significant atypical lymphocytosis which is a bad prognostic indicator according to studies.²⁸

Our study had a lower proportion of high MPV cases with leucopenia similar to study by U Ralpanowa et al²⁹ which attribute to prognostic relevance or claim good prognosis for leucopenia.²⁹

Our study is in concordance with four studies by S R Dewi et al and Manoharan A et al, which claim an association of high MPV with severe Dengue^{24,30} however other studies differ.^{15,21–23}

We had significant number of high MPV cases in association with antibody pattern. In Dengue, IgM levels rise at 3–5 days of fever and IgG at 6–15 days.^{31,32} This coincides with the time of platelet activation and recovery around 8 – 10th day.^{6,7} Our study is in concordance with several studies which associate high MPV with platelet activation and recovery.^{12,15,23,24}

6. Limitations of the Study

The study was limited by lack of

1. Standardization of MPV.
2. Baseline and serial monitoring of MPV in association with other tests.
3. Studies along with similar lines to compare data with clinical correlation.
4. Small sample size.
5. Inclusion of other platelet indices which could increase sensitivity and specificity.

7. Conclusion

Platelet indices have been investigated for their utility in Dengue. Our study on MPV suggests that it could be used as a predictor of severe dengue due to significant association with other prognostic predictors like hematocrit, thrombocytopenia and atypical lymphocytosis. A high MPV is suggestive of platelet activation and recovery and thus can impact transfusion therapy.

8. Source of Funding

None.

9. Conflict of Interest

None.

References

1. Tahlan A, Bhattacharya A. Haematological profile of dengue fever. *Int J Res Med Sci.* 2017;5(12):5367.
2. Praveen A, Shinde. Evaluation of platelet count in pediatric patients with dengue fever: a hospital based study. *Int J Contemp Med Res.* 2017;4(5):1175–7.
3. Reddy GC, Nagendra K. Clinical and diagnostic features of dengue haemorrhagic fever in children. *Int J Contemp Pediatr.* 2018;5(3):791–7.
4. Yashaswini LS, Priya. A study of hematological parameters and requirement of platelet transfusion in dengue fever. *Int J Adv Med.* 2017;4(6):1668.
5. Sreenivasa B, Manjunatha B, Nivil J. Bleeding manifestations in dengue and their correlation with the platelet count. *Sri Lanka J Child Health.* 2017;46(3):218.
6. Mohamed B. Mean platelet volume a marker for bleeding risk; a prospective study among Sudanese patients with dengue virus infection. *Sudan Med J.* 2018;p. 54.
7. Ojha A, Nandi D, Batra H. Platelet activation determines the severity of thrombocytopenia in dengue infection. *Sci Rep.* 2017;7.
8. Lee TH, Wong JGX, Leo YS, Thein TL, Ng EL, Lee LK. Potential Harm of Prophylactic Platelet Transfusion in Adult Dengue Patients. *PLoS Negl Trop Dis.* 2016;10(3):4576.
9. Ali M, Sharma A. Prophylactic platelet transfusion in dengue: Blanket of false comfort? *Natl Med J India.* 2018;31(1):24.
10. Khairkar PS, More S, Pandey A, Pandey M. Role of mean platelet volume (MPV) in diagnosing categories of thrombocytopenia. *Indian J Pathol Oncol.* 2016;3(4):606.
11. Mechin SJ, Bugg C. MPV- a quick, easy determinant of thrombotic risk. *J Thromb Hemostasis.* 2010;8:146–7.
12. Khandal A, Kumar. MPV and DENGUE. *Int J Innov Res Adv Stud.* 2017;4(6).
13. Gulati I, Kumar H, Sheth J, Dey I. Diagnostic implication of mean platelet volume in thrombocytopenia. *Med J Dr DY Patil Univ.* 2017;10(4):370.
14. Afsar N, Afroze IA, Humaira S, Abid Z. Use of Mean platelet Volume as an initial diagnostic marker in evaluation of dengue fever. *Ann Pathol Lab Med.* 2017;4(3):A310–3.
15. Kantharaj A. Role of red cell and platelet indices as a predictive tool for transfusions in dengue. *Glob J Transfus Med.* 2018;3(2):103.
16. Budak YU, Polat M, Huyral K. The use of platelet indices, platelet count, MPV, PDW in emergency non traumatic abdominal surgery- A systematic review. *Biochem Med (Zogreb).* 2016;26(2):176–83.
17. Korniluk A, Koper-Lenkiewicz OM, Kamińska J, Kemona H, Dymicka-Piekarska V. Mean Platelet Volume (MPV):

- New Perspectives for an Old Marker in the Course and Prognosis of Inflammatory Conditions. *Mediators Inflamm.* 2019;doi:10.1155/2019/9213074.
18. Schmoeller D, Picarelli MM, Munhoz TP, de Figueiredo CEP, Staub HL. Mean Platelet Volume and Immature Platelet Fraction in Autoimmune Disorders. *Front Med.* 2017;4:146.
 19. Gogaram, Prasad H, Misha S. Evaluation of Platelet Indices among Patients with Dengue Infection attending OBGY OPD/IPD in District Hospital, Churu, Rajasthan. *Appl Physiol Anat Dig.* 2018;(3):26–36.
 20. Kottke K, Marchant, Corcoran G. The Laboratory Diagnosis of Platelet Disorders. *Arch Pathol Lab Med.* 2002;126(2):133–46.
 21. Mukker P, Kiran S. Platelet indices evaluation in patients with dengue fever. *Int J Res Med Sci.* 2018;6(6):2054.
 22. Sharma K, Yadav. Association of MPV with Severity, Serology & Treatment Outcome in Dengue Fever: Prognostic Utility. *J Clin Diagn Res.* 2009;(11):1–03.
 23. Prakash GM, Anikethana GV. Use of MPV & PDW in Predicting Trend in Platelet Count and Bleeding Risks in Patients with Dengue Fever. *Int J Adv Med.* 2016;3(3):611–3.
 24. Dewi USR, Sukorini US, Intansari. Correlation of enhancement MPV with Severity of Dengue Infection. In: IOP Conference Series: Materials Science and Engineering. vol. 434; 2018.
 25. Kumar N. Correlation of Mean Platelet Volume and Platelet Distribution Width in risk categories of Dengue Fever-A Pilot Study. *J Evol Med Dent Sci.* 2018;7(2):142–5.
 26. Mangione JN. The association of cytokines with severe Dengue in children. *Trop Med Health.* 2014;42(4):137–44.
 27. Fan Z. Mean Platelet Volume and Platelet Distribution Width as markers in diagnosis of Acute Gangrenous Appendicitis. *Dis Markers.* 2015;150.
 28. Eu-Ahsunthornwattana N, Eu-ahsunthornwattana J, Thisyakorn U. Peripheral blood count for Dengue severity prediction: A Prospective Study in Thai Children. *Pediatr.* 2008;121(2):S127–8.
 29. Ralapanawa U, Alawattegama ATM, Gunrathne M, Tennakoon S, Kularatne SAM, Jayalath T. Value of peripheral blood count for dengue severity prediction. *BMC Res Notes.* 2018;11(1):400.
 30. Manoharan A, Sinha P. Role of mean platelet volume in dengue fever—a prognostic marker. *Int J Res Rev.* 2017;4(6):72–7.
 31. Fry RS. The diagnostic sensitivity of Dengue rapid test assay is significantly enhanced by using a combined Antigen and Antibody testing approach. *PLOS Nagaland Tropical Diseases. PLOS Nagaland Trop Dis.* 2011;5(6).
 32. Jang WS, Kwak SY, May WL, Yang DJ, Nam J, Lim CS. Comparative evaluation of three dengue duo rapid test kits to detect NS1, IgM and IgG associated with acute dengue in children in Myanmar. *PLoS One.* 2019;14(3):213451–213451.

Author biography

Anagha A Joshi Blood Bank Officer

Sathyavathi S Assistant Professor

Gayathri B R Tutor

Rukmini Tanjore Final Year Student

Krishnamurthy B N Associate Professor

Cite this article: Joshi AA, Sathyavathi S, Gayathri B R, Tanjore R, Krishnamurthy B N. Mean platelet volume: Has it the capacity to predict the concentration of Platelet counts in Dengue: A study. *Indian J Pathol Oncol* 2020;7(3):352–356.