Case Report

Sarcomatoid squamous cell carcinoma of urinary bladder

S V R Raja Sekhar^{1,*}, B Ramana², M Shamili¹, B Gouthami¹, Ch. Asha Latha¹, G Harshitha¹¹Dept. of Pathology, GEMS & Hospital, Srikakulam, Andhra Pradesh, India²Dept. of Urology, GEMS & Hospital, Srikakulam, Andhra Pradesh, India

ARTICLE INFO

Article history:

Received 08-01-2020

Accepted 14-02-2020

Available online 19-08-2020

Keywords:

Sarcomatoid carcinoma

Urinary bladder

ABSTRACT

Among the urinary bladder malignancies, infiltrating urothelial carcinoma is the most common while sarcomatoid variant is a rare malignancy. As the literature shows only few published case reports, We present a case of polypoidal mass of urinary bladder which was excised by TURBT (Transurethral Resection of Bladder Tumor) procedure and histopathologically diagnosed as sarcomatoid squamous cell carcinoma.

© 2020 Published by Innovative Publication. This is an open access article under the CC BY-NC license (<https://creativecommons.org/licenses/by-nc/4.0/>)

1. Introduction

According to the WHO classification of urinary bladder, Urothelial carcinoma is the most common bladder neoplasm.¹ Sarcomatoid carcinoma is a rare variant of bladder carcinomas with aggressive behavior, accounting for only 0.1% to 0.3% of carcinomas.² These types of tumors were previously called as "carcinosarcoma" as the tumor has epithelial and mesenchymal component, but the recent WHO classification of urinary tract tumors uses the term "sarcomatoid carcinoma" for all these lesions. The epithelial component most commonly seen is urothelium, followed by squamous or small cell or adeno in nature, while sarcomatous component can be of homologous or heterologous differentiation.² We present a case of sarcomatoid carcinoma having a squamous epithelial component, which is very rare.

2. Case Report

A 65-year-old male, a farmer by occupation, complaints of dysuria for four months duration associated with hematuria and abdominal pain. Ultrasound abdomen

revealed a well distended bladder with mixed echogenic mass arising from urinary bladder posterior wall measuring 7.9x4.6cm. TURBT was done and sent for histopathological examination.

Grossly Received multiple grey tan to grey-brown soft tissue bits altogether measuring 5x4x3cm. Microscopy Showed predominantly high-grade spindle cell component (70%) with myxoid stroma and moderate to poorly differentiated squamous epithelial component in solid nests and sheets (Figure 1). These tumor cells showed marked nuclear pleomorphism. Heterologous elements were not identified.

3. Discussion

Sarcomatoid carcinoma is uncommon and aggressive tumor of urinary bladder that occurs predominantly in elderly male smokers of the seventh decade. Most patients present with hematuria, dysuria, urinary retention and urinary tract infection.³ A previous history of carcinoma treated by radiotherapy or exposure to cyclophosphamide therapy is common.³

Among the different neoplasms, sarcomatoid carcinoma has a unique aggressive pathway in its pathogenesis. The exact origin of this variant was not clearly explained till

* Corresponding author.

E-mail address: rajsanuvada@gmail.com (S. V. R. R. Sekhar).

Fig. 1: **A:** Highly pleomorphic spindle cells arranged in fascicles; **B:** Spindle cells admixed with myxoid stroma; **C:** Neoplastic squamous epithelium in solid islands; **D:** Squamous sheets with intervening myxoid stroma

date. Some authors believe that it is due to synchronous appearance of two independent monoclonal neoplasms, whereas others suggest that they originate from a single neoplastic cell but shows epithelial and mesenchymal differentiation, which was also supported by recent molecular studies.⁴

Urinary bladder lateral walls are the common sites of its origin. Grossly they have a red fungating growth into the lumen of the bladder. Microscopically the tumor shows divergent differentiation with epithelial and mesenchymal components.⁵ Majority of the areas in these lesions show mesenchymal differentiation having high grade spindle cells in most of these tumors. The epithelial component is usually in the form of invasive urothelial carcinoma. The other uncommon epithelial components being squamous cell carcinoma, adenocarcinoma, or small cell carcinoma. While, osteosarcomatous component is the most common heterologous differentiation, others being chondrosarcomatous, rhabdomyosarcomatous, leiomyosarcomatous,

liposarcomatous and angiosarcomatous elements but does not have prognostic significance.^{6,7} Immunohistochemistry is useful in difficult cases.

The poor prognosis of carcinosarcoma is because of advanced stage of the disease at the time of presentation and high-histological grade and this is usually worse than prognosis of high grade invasive urothelial carcinoma.^{8,9} The appropriate standard treatment has not yet been defined because of its rarity; however, the aggressive behavior suggests radical therapy whenever possible.⁸ Total cystectomy, often followed by radiation therapy and/or chemotherapy, seems to be the preferred treatment.¹⁰ The effectiveness of these treatments is not known because of the varying results of each case. The factors predictive of long-term survival at the time of presentation are negative surgical margins and the absence of metastatic disease.⁹ Unfortunately, cases with metastasis have a very poor prognosis. Studies with a larger number of cases are needed to establish standard protocols for the treatment of sarcomatoid carcinoma of the urinary bladder

4. Conclusion

Sarcomatoid carcinomas are rare and aggressive neoplasms of urinary bladder, having two malignant components; epithelial and mesenchymal differentiation, with a note on heterologous elements. As this disease has very poor prognosis, immediate diagnosis, and treatment help in increasing the survival period for the patients.

5. Source of Funding

None.

6. Conflicts of Interest

None declared.

References

1. Zhai QJ, Black J, Ayala AG, Ro JY. Histologic variants of infiltrating urothelial carcinoma. *Arch Pathol Lab Med.* 2007;131:1244–56.
2. Lopez-Beltran A, Pacelli A, Rothenberg HJ, Wollan PC, Zincke H, Blute ML, et al. Carcinosarcoma and sarcomatoid carcinoma of the bladder: Clinicopathological study of 41 cases. *J Urol.* 1998;159(5):1497–1503.
3. Wang J, Wang FW, LaGrange CA, III GPH, Kessinger A. Clinical Features of Sarcomatoid Carcinoma (Carcinosarcoma) of the Urinary Bladder: Analysis of 221 Cases. *Sarcoma.* 2010;2010. doi:10.1155/2010/454792.
4. Cheng L, Zhang S, Alexander R, MacLennan GT, Hodges KB, Harrison BT, et al. Sarcomatoid carcinoma of the urinary bladder: the final common pathway of urothelial carcinoma dedifferentiation. *Am J Surg Pathol.* 2011;35:34–46.
5. Dahm P, Gschwend JE. Malignant Non-Urothelial Neoplasms of the Urinary Bladder: A Review. *Eur Urol.* 2003;44(6):672–81.
6. Lopez-Beltran A, Cheng L. Histologic variants of urothelial carcinoma: differential diagnosis and clinical implications. *Hum*

Pathol. 2006;37:1371–88.

7. Cheng L, Lopez-Beltran A, Bostwick DG. Sarcomatoid carcinoma (carcinosarcoma). In: *Bladder Pathology.* vol. 58. Hoboken, New Jersey: Wiley-Blackwell; 2012. p. 355–81.
8. Black PC, Brown GA, Dinney CPN. The impact of variant histology on the outcome of bladder cancer treated with curative intent. *Urol Oncol.* 2009;27(1):3–7.
9. Wright JL, Black PC, Brown GA, Porter MP, Kamat AM, Dinney CP, et al. Differences in Survival Among Patients With Sarcomatoid Carcinoma, Carcinosarcoma and Urothelial Carcinoma of the Bladder. *J Urol.* 2007;178(6):2302–6.
10. Baseskioglu B, Duman BB, Kara IO, Can C, Yildirim M, Acikalin M. Early Detection and Gemcitabine/Cisplatin Combination Positively Effect Survival in Sarcomatoid Carcinoma of the Urinary Bladder. *Asian Pac J Cancer Prev.* 2012;13(11):5729–33.

Author biography

S V R Raja Sekhar Assistant Professor

B Ramana Assistant Professor

M Shamili Assistant Professor

B Gouthami Assistant Professor

Ch. Asha Latha Assistant Professor

G Harshitha Postgraduate

Cite this article: Sekhar SVRR, Ramana B, Shamili M, Gouthami B, Latha CA, Harshitha G. Sarcomatoid squamous cell carcinoma of urinary bladder. *Indian J Pathol Oncol* 2020;7(3):505-507.