Content available at: <https://www.ipinnovative.com/open-access-journals>

Indian Journal of Pathology and Oncology

Journal homepage: www.ijpo.co.in

Case Report

Malignant mixed germ cell tumours- A rare case report with unusual presentation in a 05 years old female child

Ritu Sonkar¹, Harendra Kumar^{1,*}, Pooja Nagayach¹, Puneet Srivastava²

¹Dept. of Pathology, Sarojini Naidu Medical College, Agra, Uttar Pradesh, India

²Dept. of Surgery, Sarojini Naidu Medical College, Agra, Uttar Pradesh, India

ARTICLE INFO

Article history:

Received 22-04-2021

Accepted 23-06-2021

Available online 12-08-2021

Keywords:

MGCT

Yolk sac tumour

AFP

ABSTRACT

Mixed germ cell tumours (MGCT) of the ovary are malignant neoplasms of the ovary comprising of two or more types of germ cell components. Most of the malignant MGCT consists of dysgerminoma accompanied by endodermal sinus tumours, immature teratoma or choriocarcinoma. There are only few case reports of MGCTs with different combinations of malignant components. We present a very rare case of malignant MGCT in a 05 years old female child, who presented with difficulty in defecation for 6 months, swelling over sacral region for 2 months, along with bilateral inguinal swellings and difficulty in urination for 1 week. Swelling was firm, non-mobile & non-tender measuring 6.6x5 cm. On cytology it was diagnosed as malignant germ cell tumour, which was confirmed on histology as malignant MGCT (predominantly yolk sac tumour >90% and embryonal carcinoma (<10%)), metastasizing into bilateral inguinal lymph nodes.

This is an Open Access (OA) journal, and articles are distributed under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License](https://creativecommons.org/licenses/by-nc-sa/4.0/), which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: reprint@ipinnovative.com

1. Introduction

Globally ovarian cancer accounts for 2,25,000 new cases and 1,40,000 deaths occurs annually.¹ Malignant germ cell tumours (GCT) of ovary, an uncommon neoplasm originates from the primitive germ cells of embryonic gonad. Mostly they occur in the second and third decade of life and accounts for 2–3% of all ovarian neoplasms.² The survival rates for germ cell tumours have dramatically improved for the past three decades coincident with more aggressive surgical staging and combination therapy.³ World Health Organization (1973) has classified malignant germ cell tumours as endodermal sinus tumour (yolk sac tumour), immature teratoma and dysgerminoma, non-gestational choriocarcinoma, embryonal carcinoma, and mixed germ cell type.⁴ Malignant mixed germ cell tumour is a type of tumour which consists of two or more malignant germ cell components. These tumours are rare cancers, which are seen

in 8% cases of germ cell tumours but are very aggressive in nature.⁵ The Endodermal Sinus Tumour (EST) and dysgerminoma are the most common combination reported in the literature⁶ and the embryonal carcinoma and immature teratoma are the rarest combination as its components.⁴ Embryonal carcinoma, although very rare, but it holds the very malignant potential.⁷ In review of literatures, teratoma is the most common histology encountered followed by the dysgerminoma.³

2. Case Report

A 5-years old female child presented to the surgery department with a brief history of difficulty in defecation for 6 months, swelling over sacral region for 2 months, along with bilateral inguinal swellings and difficulty in urination for 1 week. On physical examination swelling was firm, non-mobile & non-tender measuring 6.6x5 cm (Figure 1 a & 1b). Complete blood counts, Liver function tests and Kidney function tests were within normal limit. Ultrasound was

* Corresponding author.

E-mail address: dr_haren@yahoo.co.in (H. Kumar).

suggestive of heterogenous mass in sacral region, bilateral inguinal lymphadenopathy. MRI findings revealed a large solid presacral and pre-coccygeal mass lesion with pelvic extension, inguinal lymphadenopathy with mass effect and extension, suggestive of sacrococcygeal teratoma (Type III). Serum alfa fetoprotein (AFP) level was 39773.80 ng/ml. FNAC from sacral mass & bilateral inguinal swelling revealed high cellularity, comprising of neoplastic cells arranged in solid sheets, papillaroid fragments loose clusters and scattered singly. Individual tumour cells have high N:C ratio, with oval to elongated nuclei with coarse chromatin, moderate amount of vacuolated cytoplasm. At places eosinophilic material is also seen within the cell clusters (Figure 2 a & b). Cytological features were suggestive of malignant germ cell tumour of ovary. Excisional biopsy was done and sent for histopathological examination to our department. On gross examination, multiple greyish white pieces of tissue were received, altogether measuring 4x3x1 cm. Microscopically, multiple sections examined showed tumour arranged in reticular pattern. These tumour cells showed vesicular nuclei with moderate pleomorphism (Figure 3 c). Perivascular arrangement of tumour cells was also seen (Schiller-Duval bodies) (Figure 3 d). Some areas showed papillary and glandular differentiation (Figure 1 a). Some areas also showed undifferentiated primitive cells having hyperchromatic nuclei with coarse chromatin and occasional giant cells. Small focus of necrosis also noted (Figure 3 b). Lymphoid tissue was also identified infiltrated by tumour cells as described above. IHC stain shows cytoplasmic positivity for AFP and Pancytokeratin (Figure 4 a & b). Histopathological features were suggestive of malignant mixed germ cell tumour (predominantly yolk sac tumour (>90%) and embryonal carcinoma (<10%), metastasizing into the inguinal lymph nodes. Patient was treated with combination chemotherapy (cisplatin, etoposide and bleomycin) and she responded to the treatment. Then she was planned for debulking surgery.

Fig. 1: Sacral swelling with skin discoloration (a): along with bilateral inguinal swellings (b)

Fig. 2: Cellular smears showing solid sheets, papillaroid fragment, loose clusters and singly scattered cells (FNAC, MGG 100X (a), 400X (b))

Fig. 3: Section examined showing papillary and glandular differentiation along with foci of necrosis (H&E, 40x) (a), Embryonal component showing primitive cells and necrotic area, (H&E, 400x) (b), Yolk sac component showing reticular areas (H&E, 100x) (c), Schiller-Duval body (H&E, 100x) (d)

Fig. 4: a: AFP positivity; b: Cytokeratin positivity (IHC, 400X)

3. Discussion

The rare combination of uncommon germ cell components, large sacral mass with bilateral inguinal lymphadenopathy, and very high levels of tumour markers makes this case unusual. The germ cell tumour generally present at an average age of 13.8 years (4–27 years).⁸ Most commonly patient presents with complaint of abdominal mass which may be associated with or without abdominal pain or fever. Embryonal carcinoma can also present with precocious

puberty or irregular vaginal bleeding as it may secrete estrogen.⁹ Immunohistochemistry is a widely used for the confirmation of diagnosis and to find new management modalities.^{10,11} Yolk sac tumour shows positivity for AFP, Glypican-3, SALL4 and LIN 28, whereas in embryonal carcinoma tumour cells are positive for CK (AE1/AE3), CD30, OCT4, SALL4 and Glypican-3.¹² Trinh DT et al. has described the utility of tumour markers CD117, CD133, SALL4, OCT4, TCL1 and glypican-3 in the malignant germ cell tumours of ovary.¹¹ Ovarian endodermal sinus tumours are highly aggressive but with surgery and combination chemotherapy, the five-year survival with stage I tumours and more advanced disease are 92% and 29% to 44% respectively.¹³ Survival in embryonal carcinoma the first reported series of 15 patients was 39%, with 50% of stage I patients being disease free at 3.75 to 15 years post-surgery and chemotherapy.¹⁴ More recent data illustrate the conservative surgery and combination chemotherapy improves the survival, with survival rates of 98% and 94% for early and advanced stage tumours.¹⁵ Elevation of both AFP and hCG levels is a strong predictor of poor outcome.¹⁶ Kumar et al. suggested lymph node involvement as an independent predictor of poor survival.¹⁷ Pectasides et al. declared that if tumour appears to be confined to ovaries or pelvis, a pelvic and para-aortic lymphadenectomy should also be performed. If upper abdominal involvement is there, an effort to remove all visible tumour should be made. Also, in view of the excellent chemosensitivity, clinical judgment can be used if surgical management would significantly increase postoperative morbidity.¹⁸ If infertility is concerned than unilateral oophorectomy and surgical staging is the minimal surgery in ovarian germ cell tumour. Preservation of fertility is recommended even in metastatic disease as these tumours are highly sensitive to chemotherapy. Outcome cannot be changed by hysterectomy and bilateral salpingo-oophorectomy.¹⁰

4. Conclusion

Ovarian germ cell tumours arise from primordial germ cell derived from the embryonal gonads. Debulking surgery and combination chemotherapy is the mainstay in treatment of ovarian cancer.

5. Conflicts of Interest

All contributing authors declare no conflicts of interest.

6. Source of Funding

None.

References

1. Ferlay J, Shin HR, Bray F, Forman D, Mathers C, Parkin DM. Estimates of worldwide burden of cancer in 2008: GLOBOCAN. *Int*

J Cancer. 2008;127(12):2893–917.

2. Quirk JT, Natarajan N. Ovarian cancer incidence in the United States, 1992–1999. *Gynecol Oncol*. 2005;97(2):519–23. doi:10.1016/j.ygyno.2005.02.007.
3. Smith HO, Berwick M, Verschraegen CF, Wiggins C, Lansing L, Muller CY, Qualis CR (2006) Incidence and survival rates for female malignant germ cell tumors. *Obstet Gynecol*. 2005;107(5):1075–85.
4. Koshy M, Vijayanathan A, Vadiveloo V. Malignant ovarian mixed germ cell tumour: a rare combination. *Biomed Imaging Interv J*. 2005;1(2):e10.
5. Munemane A, Munemane M. Malignant mixed germ cell tumours of ovary. A Report of two cases. *Int J Med Sci and Public Health*. 2014;3(1):105–7. doi:10.5455/ijmsph.2013.061020133.
6. Tiwary B, Sinha H, Pandey V. Malignant mixed germ cell tumor of ovary: a rare case report. *Int J Reprod Contracept Obstet Gynecol*. 2015;4(2):511–3. doi:10.5455/2320-1770.ijrcog20150450.
7. Moniaga NC, Randall LM. Malignant Mixed Ovarian Germ Cell Tumor with Embryonal Component. *J Pediatr Adolesc Gynecol*. 2011;24(1):e1–e3. doi:10.1016/j.jpap.2010.05.001.
8. Brammer HM, Buck JL, Hayes WS, Sheth S, Tavassoli FA. From the archives of the AFIP. Malignant germ cell tumors of the ovary: radiologic-pathologic correlation. *RadioGraphics*. 1990;10(4):715–24. doi:10.1148/radiographics.10.4.2165627.
9. Berek JS, Hacker N, Berek JS, Hacker NF. Practical Gynecologic Oncology. 5th ed. Philadelphia: Williams and Wilkins; 2007.
10. re JBJ, Ali NB, Fadhlou A, Dubuisson JB, Guillou L, Sutter A, et al. Two Case Reports of a Malignant Germ Cell Tumor of Ovary and a Granulosa Cell Tumor: Interest of Tumor Immunohistochemistry in the Identification and Management. *Front Oncol*. 2014;4. doi:10.3389/fonc.2014.00097.
11. Trinh DT, Shibata K, Hirose T, Umezaki T, Mizuno M, Kajiyama H, et al. Diagnostic utility of CD117, CD133, SALL4, OCT4, TCL1 and glypican-3 in malignant germ cell tumors of the ovary. *J Obstet Gynaecol Res*. 2012;38(5):841–8. doi:10.1111/j.1447-0756.2011.01798.x.
12. Fletcher CDM. Diagnostic histopathology of tumours. vol. 1. 3rd ed.; 1995. p. 604–15.
13. Kawai M, Kano T, Furuhashi Y, Misawa T, Nakashima N, Hattori S. Prognostic factors in yolk sac tumors of the ovary. A clinicopathologic analysis of 29 cases. *Cancer*. 1991;67(1):184–92.
14. Kurman RJ, Norris HJ. Embryonal carcinoma of the ovary. A clinicopathologic entity distinct from endodermal sinus tumor resembling embryonal carcinoma of the adult testis. *Cancer*. 1976;38(6):2420–33. doi:10.1002/1097-0142(197612)38:6<2420::aid-cncr2820380630>3.0.co;2-2.
15. Low JH, Perrin LC, Crandon AJ, Hacker NF. Conservative surgery to preserve ovarian function in patients with malignant ovarian germ cell tumors. *Cancer*. 2000;89(2):391–8. doi:10.1002/1097-0142(20000715)89:2<391::aid-cncr26>3.0.co;2-v.
16. Murugesu N, Schmid P, Dancey G, Agarwal R, Holden L, McNeish I, et al. Malignant Ovarian Germ Cell Tumors: Identification of Novel Prognostic Markers and Long-Term Outcome After Multimodality Treatment. *J Clin Oncol*. 2006;24(30):4862–6. doi:10.1200/jco.2006.06.2489.
17. Kumar S, Shah JP, Bryant CS, Imudia AN, Cote ML, Ali-fehmi R, et al. The prevalence and prognostic impact of lymph node metastasis in malignant germ cell tumors of the ovary. *Gynecol Oncol*. 2008;110(2):125–32. doi:10.1016/j.ygyno.2008.04.022.
18. Pectasides D, Pectasides E, Kassanos D. Germ cell tumors of the ovary. *Cancer Treat Rev*. 2008;34(5):427–41. doi:10.1016/j.ctrv.2008.02.002.

Author biography

Ritu Sonkar, Junior Resident

Harendra Kumar, Associate Professor

Pooja Nagayach, Assistant Professor

Puneet Srivastava, Associate Professor

Cite this article: Sonkar R, Kumar H, Nagayach P, Srivastava P. Malignant mixed germ cell tumours- A rare case report with unusual presentation in a 05 years old female child. *Indian J Pathol Oncol* 2021;8(3):416-419.