Content available at: <https://www.ipinnovative.com/open-access-journals>

Indian Journal of Pathology and Oncology

Journal homepage: www.ijpo.co.in

Original Research Article

Correlation of ER PR receptors with other prognostic parameters of breast carcinoma

Nekta Anand¹, Vishrabdha Rahul Pawar^{2,*}, Shahnaz Zaidi¹¹Dept. of Pathology, Bharati Vidyapeeth (Deemed to be University) Medical College & Hospital, Sangli, Maharashtra, India²Dept. of Pathology, Dr. D Y Patil Medical College, Kolhapur, Maharashtra, India

ARTICLE INFO

Article history:

Received 03-07-2021

Accepted 14-08-2021

Available online 23-11-2021

Keywords:

Breast carcinoma

Estrogen and progesterone receptor

ABSTRACT

Aim & Objectives: To study the MRM specimens for size, grade of the tumor, LVI and LN metastases and the ER/PR receptors of the tumor. To study the association of ER/PR status with the above mentioned prognostic parameters.

Materials and Methods: This three-year study includes 72 histopathologically confirmed cases of carcinoma breast. The tumor type, grade, LNM, LVE were reported on H & E. The ER-PR study was done of all 72 cases. The tumor size, grade, LNM, LVE were correlated with receptor status.

Observation: Out of 72 cases majority (53%) of cases were in 41-50 years of age group, and tumor size was between 2-4 cm in (54.16%) cases; majority were of grade II carcinoma (60%) of which 59% were ER-PR positive. Out of 46 cases with negative LNM and LVE, 30 cases (65%) were ER-PR positive.

Conclusion: In our observation there is no association between higher histological grade and ER-PR status. No obvious correlation with tumor size was noted. But increase in tumor size could also be a poor predictor of ER-PR status. LNM, LVE is poor predictor of ER-PR status.

This is an Open Access (OA) journal, and articles are distributed under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License](https://creativecommons.org/licenses/by-nc-sa/4.0/), which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: reprint@ipinnovative.com

1. Introduction

Breast cancer is the most frequent cancer in females in the world and is the second most common cancer in India after cervical cancer. The breast cancer is second most common reason of loss of existence in each developed and additionally most of the growing countries.¹ For the appraisal of prognosis of breast cancer commonly followed is the Nottingham Prognostic Index (NPI) which includes-tumor size, histological tumor grade, lymph node metastasis and hormone receptor status. These prognostic component help in administration and therapeutic requirement of breast cancer patient.²

Technological advance has extended the perception of the molecular groundwork of tumor prognosis and its

treatment response. Over the last 10 years, these advances have led to identification of biomarker.³ Hormone receptor frequently estrogen receptor (ER), Progesterone receptor (PR) and human epidermal receptor (HER- 2) are most applicable scientific biomarker which is superbly used in case of identification and selection of specific therapy for the breast cancer. Estrogen and progesterone receptor performs a very necessary attribute in the treatment of breast cancer.⁴ The progesterone receptor acts as modulator of estrogen expression and function. Progesterone receptor expression lengthens along with atypia of breast tissue.⁵

In this study we have studied the correlation of ER/ PR status with lymph nodes metastasis, lymphovascular emboli and also with distant metastasis.

ER is the classic and most recounted biomarker that is prevalent in breast tumor. The PR or nuclear receptor subfamily 3, group C, member 3 is a protein present inside

* Corresponding author.

E-mail address: vishrabdhapawar65@gmail.com (V. R. Pawar).

the cell.⁴

Since the breast cancer is a heterogeneous disease, so it is important to decide the ER/PR and HER-2/Neu status in lymph node metastasis, lymphovascular emboli and distant metastasis.⁶

The decisiveness of ER and PR response in breast cancer is ameliorated medical practices now- a- days, with the useful resource of which it is very convenient to predict the response of the patients. ER status is critical in prediction of the response to adjuvant Tamoxifen therapy, while PR is an ER modulator and additionally an estrogen regulated gene. Simultaneous analysis of ER, PR is more informative regarding the prediction of likely hormonal response because more than half of ER positive tumors are positive with PR expressions.⁷

Endocrine hormonal therapy for first breast cancers was first started in late 1980 by BEATSON. He located the response in few young females with enhanced breast cancer after he performed oophorectomy. Subsequent scientific studies suggest the effectiveness of endocrine treatment on breast cancer patients which reduces the morbidity and mortality. The use of estradiol as treatment has been observed to have a positive impact on premenopausal females. Tamoxifen was initially used as antitestosterone but later was found to be an effective treatment of breast cancer. So now it comes under category of SERM. Although endocrine remedies are higher tolerated than chemotherapy but may additionally have problems occasional complications like thromboembolic events, uterine cancer and some agent are much less compatible to the patients. Similarly, HER-2 receptor serve as an essential marker for advantages of treatment focused on this receptor tyrosine kinase like Trastuzumab and Lapatinib.⁶

2. Materials and Methods

This study was approved by the Institutional Ethical Committee, IEC number given was 354/19.

In this three year study we have included 72 histopathological confirmed cases of Breast carcinoma.

The MRM specimen obtained was embedded in 10% formalin, along with the requisition form which included the clinical data of the patient. Grossing of the specimen was done to note down the size of the tumor and other routine details. Apart from those sent by the surgeon the lymph nodes were meticulously dissected in the axillary tail.

The necessary bits were given to prepare the paraffin tissue blocks. Hematoxylin and eosin staining was done on the slides. The slides were examined to confirm and type the breast carcinoma. For ER and PR study the sections were cut at 4 micrometer thickness and mounted onto a slide and left to dry overnight at 37°C. Section were deparaffinized and rehydrated. Antigen retrieval was achieved by heat retrieval using a bench autoclave. Briefly slides were placed in coplin jars containing enough

Tris EDTA (pH-9.0) to cover the sections, then autoclave at 121°C for 15 min for both ER and PR. After washing, the sections were covered by applying 3% H₂O₂ and the slides were then rinsed with phosphate buffer solution (PBS). Slides were incubated with 100-200 microliter of primary antibodies for 30 min. at 37°C temperature in a moisture chamber, then rinsed in PBS.

Finally, the section was washed 4 times for 4 minutes with changes of phosphate buffer solution, followed by adding 3,3', diaminobenzene tetra hydrochloride as chromogen to produce the characteristic brown stain. The sections were counterstained, dehydrated and mounted for analysis for each run of staining, a positive control slide was also prepared. The positive control slides were prepared from breast carcinoma known to be positive for the antigen under study. The negative control slides were prepared from the same tissue block, but incubated with PBS instead of primary antibody. The tumor size was assessed by specimen. Tumor less than 2 cm taken as T1, tumor size 2-4 cm were taken as T2 and tumor more than 4 cm were taken as T3. Under the Nottingham combined histological system was used for grading. grade 1 carcinoma includes tumor with combined score of 3,4 or 5; Grade 2 includes score of 6 and 7; Grade 3 includes tumor with score of 8 and 9. Mitotic index was assessed. Each risk factor was compared with receptor status.

3. Results

The result of our study showed positive correlation between ER/PR with low grade tumor.

These findings showed the importance of the biomarkers because they provide valuable prognostic information for best treatment decision. Breast carcinoma is the common malignant tumor.

53% cases were between 40-60 year of age and majority (54%) tumors were of 2-4 cm in size. 39% tumors were with grade II carcinoma. The findings in our studies show the negative correlation of ER-PR marker with increasing tumor grade and lymphovascular embolization and lymph node metastasis. The immunohistochemical marker have the prognostic and predictive values.

Table 1: Distribution of total number of cases on the basis of age group

Age groups in yrs	No. of cases	Percentage of cases
30-40	13	18%
41-50	24	33%
51-60	15	20%
>61	20	27%

Table 2: Distribution of total number of cases on the basis of age group

Age groups in yrs.	ER+	ER-	PR+	PR-
30-40	10	4	8	6
41-50	15	9	15	9
51-60	10	4	8	6
>61	17	3	12	8

Table 3: Distribution of total number of cases on the basis of age group

	No of cases	Percentage (%)	ER+PR+ (n)	(%)	ER+PR- (n)	(%)	ER-PR+ (n)	(%)	ER-PR- (n)	(%)
Tumor Size	< 2cm	04	5.55	04	100	00	00	00	00	00
	2-4 cm	39	54.16	22	56	08	11.11	00	00	09
	> 4 cm	29	40.27	16	55	00	00	01	1.38	11

Table 4: Correlation of ER/PR status with tumor grade

	No of cases	Percentage (%)	ER+PR+ (n)	Percentage (%)	ER+PR- (n)	ER-PR+ (n)	ER-PR- (n)
Tumor Grade I	24	10	16	66	03	01	05
II	43	39	24	59	05	00	13
III	05	6.94	02	40	00	00	03

Table 5: Correlation of ER/PR status with LVE/ LN metastasis/ Distance metastasis

	No of cases	Percentage (%)	ER+PR+ (n)	(%)	ER+PR- (n)	ER-PR+ (n)	ER-PR- (n)
Lymphovascular Emboli	+	26	36.11	16	61.53	01	00
	-	46	63.88	26	56.52	07	02
Lymph Node Metastasis	+	26	36.11	12	28.57	03	01
	-	46	63.88	30	71.42	05	01

4. Discussion

Breast cancer is most common tumor in women. It is highly curable if diagnosis is done at an early stage because significant number of cancer patients with early stage of carcinoma possesses microscopic metastasis at the time of diagnosis. For assessing the prognosis and management of the breast cancer ER and PR both markers work as prognostic as well as predictive factors. Correlation of expression of estrogen and progesterone with each other and to various clinicopathological parameters is very significant.⁵ Estrogen and progesterone status are measured by chemical ligand binding assay are the only prognostic and predictive biomarker recommend for routine clinical use in breast cancer by American pathologist. Both are relatively weak prognostic factor, but strong predictive factor for response to the adjuvant and therapeutic hormonal therapy.

In this article we have studied 72 MRM specimens for size, grade of the tumor, lymphovascular emboli, lymph node metastasis and ER and PR status of the tumor. We have studied the association of ER and PR status with the above mentioned parameters.

We studied 72 MRM specimens, where the breast carcinoma was common in the age group >40 years (33%). Grade II was the commonest grade amongst the 72 cases (60%), and 54.16 % the tumor size was between 2-4 cm. Similar was the observation was Premchand, Ashok Kumar et al.⁵ who studied 100 cases and majority (96.82%) of the breast carcinoma was in the age group of >40 years. 43% cases were of grade II tumor and, in 47% cases the tumor size was between 2-5 c.m. According to this, as the size and grade of the tumor increases the ER and PR positive status decreases. Our findings are similar to Mohammad Golam Mostafa where higher histologic grade is poor predictor of hormone receptor status.⁸

Fig. 1: 10x H&E- IDC-Grade I -Glandular and tubular differentiation, mild nuclear pleomorphism

Fig. 3: 40x H&E- IDC- Grade III -Sheets and nests of cells with marked nuclear atypia and mitotic activity

Fig. 2: 40x H&E- IDC-Grade II -Tubular and glandular differentiation and nests of cells, with moderately atypical nuclei, mild mitotic activity

Fig. 4: 10x -H&E- Lymph node metastasis-breast tumor metastasis in subcapsular sinus

Correlation of age of the patient with ER and PR status with other studies done. In our study 32.29% cases were of 60 years of age group and 79-80% are of >40 years of age group were ER positive, while 81% in age group of >40 were positive of PR marker.

Premchand, Ashok Kumar et al.⁵ shows similar result like our study which has majority of ER positive (96.82%) cases were of age >40 year and 44.44% were of age >60 years. The majority of PR positive (96.55%) cases were of age >40 years and 43.10% were of age >60 years. ER-PR positivity more in >40 years. ER-PR more in the tumor size (2-5) cm and in grade II and III.

Ramandeep Kaur et al.⁹ found 66% cases in <50 years with 43% of grade II carcinomas and, in 43% cases tumor size between 2-5 c.m. with significant correlation between age of the patient and ER, PR expression. ER- PR expression correlation with age at presentation more positive in postmenopausal age. No significant correlation between axillary lymph nodes status with ER, PR expression. There is a significant correlation between tumor size and tumor grade with ER, PR in above mentioned study. Grade II tumor with positive ER-PR expression. Sepideh Siadati et al.¹⁰ studied 33 cases with 34% case in the age group of >40 year, 80. 66% cases of grade II and 58.33% cases the tumor size was 2-5cm. No correlation with

Fig. 5: 40x H&E- Lymph node metastasis-breast tumor metastasis in subcapsular sinus

Fig. 7: 4X-Estrogen positive - 6/8 (weakly positive)

Fig. 6: 40x H&E- Lymphovascular emboli -breast tumor cells in the vascular space with destruction of endothelial lining of vessel

Fig. 8: 40X: Estrogen positive 6/8 (weakly positive)

axillary lymph nodes with ER, PR. No correlation with age and size. ER-PR positive status with grade I and II.

Mohammad Golam Mostafa et al.⁸ had a large sample size of 1042 cases and in 63% cases the tumor was of grade III, the predominantly affected age group was similar showing no lymphovascular emboli and lymph node metastasis. The ER and PR positive status increases with increases in tumor size.

Premchand, Ashok Kumar et al.⁵ shows similar result with negative correlation with lymphovascular emboli. The result with similar with Asmeron Tesfamarian Sengal et al.¹

and there was significant associated between age tumor size and grade, but no associated with lymph nodes. Patient with grade III and young age are more likely to be hormone receptor negative.

Manuneethimaren Thiagarajan et al.¹¹ and Mohammad Golam Mostafa et al.¹² did not find any significant correlation with ER and PR status lymphovascular emboli. ER -PR negativity increased with increase tumor size. ER negative increased with increasing tumor grade.

According to present study negative correlation is noted in lymphovascular emboli, lymph node metastasis with ER and PR status.

Fig. 9: 10X: Estrogen positive 8/8 (strongly positive)

Fig. 10: 40X: Estrogen positive 8/8 (strongly positive)

We found 63.88% cases with no lymphovascular emboli and they had positive ER and PR receptors, while other authors did not observe any significant correlation between lymphovascular emboli and ER and PR status.

5. Source of Funding

None.

6. Conflict of Interest

The authors declare no conflict of interest.

7. Acknowledgement

I would like to extend my gratitude to HOD and Professor of department of surgery Dr. Vinod Prabhu for the cases of MRM surgeries. I would like to thank to technicians of my department who helped me for this project. Secondly I would also like to thank my parents and friends who helped

me a lot in finishing this project within limited time.

References

1. Sengal AT, Haj-Mukhtar NS, Elhaj AM, Bedri S, Kantelhardt EJ, Mohamedani A. Immunohistochemistry defined subtypes of breast cancer in 678 Sudanese and Eritrean women: hospitals based case series. *BMC Cancer*. 2017;17:804. doi:10.1186/s12885-017-3805-4.
2. AlZaman AS, Mughal SA, AlZaman YS, AlZaman ES. Correlation between hormone receptor status and age, and its prognostic implications in breast cancer patients in Bahrain. *Saudi Med J*. 2016;37(1):37–42. doi:10.15537/smj.2016.1.13016.
3. Mehta S, Shelling A, Muthukaruppan A, Lasham A, Blenkiron C. Predictive and prognostic molecular markers for cancer medicine. *Ther Adv Med Oncol*. 2010;2(2):125–48. doi:10.1177/1758834009360519.
4. Shaikh F, Jamal Q, Baig S, Hadi NI, Majeed N. Correlation of hormone receptor and HER-2/neu expression with clinicopathologic parameters in primary breast tumors. *Asian Pac J Cancer Prev*. 2016;17(7):3363–7.
5. Chand P, Kumar A, Singla V, Singh G, Ramandeep R. Immunohistochemical profile of infiltrating ductal carcinoma breast- Prognostic and therapeutic use. *J Evol Med Dent Sci*. 2017;6(21):1695–1700. doi:10.14260/Jemds/2017/373.
6. Desouki MM, Atta IS, Wolff DJ, Self SE. Comparison Between HER2, Estrogen Receptors and Progesterone Receptors in Primary Breast Carcinomas and Matched Lymph Node Metastases. *Turk Patoloji Derg*. 2016;32(3):178–85.
7. Hammond MEH, Hayes DF, Dowsett M, Allred DC, Hagerty KL. American Society of Clinical Oncology/College of American Pathologists guideline recommendations for immunohistochemical testing of estrogen and progesterone receptors in breast cancer (unabridged version). *Arch Pathol Lab Med*. 2010;134(7):e48–72. doi:10.5858/134.7.e48.
8. Mohsin SK, Weiss H, Havighurst T, Clark GM, Berardo M, Roanh LD, et al. Progesterone receptor by immunohistochemistry and clinical outcome in breast cancer: a validation study. *Mod Pathol*. 2004;17(12):1545–54.
9. Kaur N, Singh H, Chand P, Kundal RK. Immunohistochemical profile of infiltration ductal carcinoma breast (Not otherwise s-A study of 100 cases). *Ann Int Med Den Res*. 2017;3(3):17–21.
10. Sepideh S, Sharbatdaran M, Nikbakhsh N, Ghaemian N. Correlation of ER, PR and HER-2/Neu with other Prognostic Factors in Infiltrating Ductal Carcinoma of Breast. *Iran J Pathol*. 2015;10(3):221–6.
11. Thiagarajan M, Navrathan N, Mohanapriya T, Kumar A, Singh B. Correlation between estrogen receptor, progesterone receptor, HER-2/neu status and other prognostic factors in carcinoma breast in Indian population. *Int Surg J*. 2015;2(4):515–22.
12. Mostafa MG, Larsen MT, Love RR. Estrogen Receptor, Progesterone Receptor, and Her-2/neu Oncogene Expression in Breast Cancers Among Bangladeshi Women. *J Bangladesh Coll Phys Surg*. 2010;28(3):157–62.

Author biography

Nekta Anand, Junior Resident
<https://orcid.org/search?searchQuery=0000-0002-3964-6592>

Vishrabdha Rahul Pawar, Professor and HOD

Shahnaz Zaidi, Tutor

Cite this article: Anand N, Pawar VR, Zaidi S. Correlation of ER PR receptors with other prognostic parameters of breast carcinoma. *Indian J Pathol Oncol* 2021;8(4):437–442.